

# MMWR

## MORBIDITY AND MORTALITY WEEKLY REPORT

---

- 745 Workplace Smoking Survey — New York City  
747 Tularemia — New Jersey  
753 Multiply Resistant Shigellosis in a Day-Care Center — Texas

### *Perspectives in Disease Prevention and Health Promotion*

---

#### **Workplace Smoking Survey — New York City**

During the period May 16-23, 1986, employees of the New York City Department of Health (NYCDOH) participated in a survey regarding smoking practices and attitudes toward a workplace smoking policy. The survey was conducted to obtain baseline information for evaluating the impact of a smoking policy initiated by the mayor and scheduled to be implemented July 1, 1986. It was also intended to familiarize employees with the policy.

Questionnaires were completed by employees who volunteered to attend one of several NYCDOH meetings concerning the mayor's proposed legislation and pending executive order about smoking restrictions. Of the estimated 900 employees in the department's primary office building, 608 attended the meetings, and 496 completed the survey. Thus, 55% of the total employees and 82% of those attending the meetings completed the questionnaire. Of the respondents, 137 (28%) currently smoked cigarettes, eight (2%) smoked pipes or cigars, 333 (67%) were nonsmokers, and 18 (4%) did not answer this question. The female to male ratio of respondents was 2.5:1. Thirty-one percent of the males and 28% of the females were current smokers. Eighteen percent of the smokers smoked a pack or more of cigarettes per day. Sixteen percent of the cigarette smokers reported that they did not smoke at work. Fifty-nine percent of nonsmokers reported at least occasional exposure to tobacco smoke from others in the workplace; 56% of nonsmokers reported at least occasional exposure to tobacco smoke from the visiting public.

Regarding employee attitudes toward smoking in the workplace, 63% of all respondents (26% of smokers and 79% of nonsmokers) reported being annoyed when other employees smoked nearby. Of nonsmokers, 38% reported that, when exposed to tobacco smoke, they would like to ask smokers to stop but are hesitant to do so. Thirty-three percent of nonsmokers reported that they were able to work without noticing smoke. Twenty-nine percent reported that they try to move away when other employees smoke. Overall, 82% of the respondents (including 69% of smokers) indicated that smoking in the workplace should be either limited (65%) or banned (17%). Most respondents indicated that restricting smoking in the NYCDOH would have no adverse effect on relations among their co-workers (87%), on their job performance (94%), or on their office morale (90%). Of current smokers, 46% indicated they would quit or reduce their smoking if workplace smoking were restricted.

### Smoking Survey — Continued

Reported by KJ Denard, MS, MK Bradstock, MD, MPH, P Clarke, MPH, Div of Health Promotion, SC Joseph, MD, MPH, New York City Dept of Health; Div of Environmental Hazards and Health Effects, Center for Environmental Health, Div of Health Education, Center for Health Promotion and Education, Epidemiology Program Office, CDC.

**Editorial Note:** The control of smoking and tobacco smoke exposure in the workplace has become an important public health issue in the United States in recent years. No studies have yet quantified the nonsmoking worker's risk of lung cancer from chronic exposure to tobacco smoke in the workplace. However, numerous studies have documented that nonsmoking wives of smoking men have a risk of lung cancer that is between 14% and 34% higher than that of wives of nonsmoking men (1). In addition, it has been shown that employees exposed to sidestream tobacco smoke in the work environment are at greater risk of developing small airways dysfunction than are nonexposed employees (2). Small airways disease, which is the first pathological change seen in beginning smokers (3), may increase the risk of developing disabling chronic airways obstruction (4).

As a consequence of data such as these concerning the effects of sidestream tobacco smoke exposure, an increasing number of employers have instituted policies to control smoking in the workplace. While some policies and control measures have been adopted voluntarily, others have been required by legislative actions. There are already laws in 17 states and ordinances in at least 100 localities regulating workplace smoking (Office on Smoking and Health, unpublished data). In one recent national survey, 36% of 662 responding employers reported having established workplace smoking policies; an additional 2% planned to enact policies by the end of 1986, and 21% reported that policies were under consideration (5).

Despite the voluntary nature of the NYCDOH survey, the results are consistent with previously reported findings concerning employee knowledge, attitudes, and smoking practices in the workplace (6-10). Since this represents 55% of total NYCDOH employees, however, these results should be interpreted with caution. Smoking prevalence among respondents in this survey (28%) is similar to the estimates of national smoking prevalence (30%) (11). It is also similar to the prevalence reported for white collar workers (32% of females, 33% of males) (12) and for those in a surveyed private workplace (33%) (6).

In most surveys, the majority of respondents have approved of some limitation of smoking in the workplace (6-10). A nationwide survey commissioned by the American Lung Association (13) asked 1,540 randomly selected individuals whether companies should have a policy on smoking at work. Eighty-seven percent of all respondents—including 80% of smokers—indicated that smoking in the workplace should be limited. Surveys of employees at individual workplaces have provided similar support for smoking restrictions. For example, 74% of employees at a large health maintenance organization approved of a smoking prohibition policy 4 months after implementation of the policy (7). In another survey, 71% of all employees indicated that smoking in the immediate work area should be restricted (80% of non-smokers and 51% of smokers) (6).

Policies limiting smoking in the workplace not only protect nonsmokers from the health effects of passive smoking but also may encourage smokers to quit or reduce smoking. In one survey, 51% of the employees who smoked indicated that workplace smoking regulations might prompt them to reduce smoking or try to quit smoking completely (6). In the NYCDOH survey, the majority of participating employees were in favor of restricting smoking in the workplace.

#### References

1. National Research Council. Passive smoking. Washington, DC: National Academy Press, 1986.
2. White JR, Froeb HF. Small-airways dysfunction in nonsmokers chronically exposed to tobacco smoke. *N Engl J Med* 1980;302:720-3.

*Smoking Survey — Continued*

3. Niewoehner DE, Kleinerman J, Rice DB. Pathologic changes in the peripheral airways of young cigarette smokers. *N Engl J Med* 1974;291:755-8.
4. Ingram RH. Chronic bronchitis, emphysema and airways obstruction. In: Petersdorf RG, Adams RD, Braunwald E, Isselbacher KJ, Martin JB, Wilson JD, eds. *Harrison's principles of internal medicine*. 10th ed. New York: McGraw-Hill Book Company, 1983:1546.
5. Bureau of National Affairs, Inc. *Where there's smoke: problems and policies concerning smoking in the workplace*. Washington, DC: Bureau of National Affairs, 1986.
6. Pacific Telephone and Telegraph Company. *Employee smoking study (project no. 82-63)*. San Francisco: Pacific Telephone Company, January 1983.
7. Rosenstock IM, Stergachis A, Heaney C. Evaluation of smoking prohibition policy in a health maintenance organization. *Am J Public Health* 1986;76:1014-5.
8. Martin MJ, Silverman MF. The San Francisco experience with regulation of smoking in the workplace: the first twelve months. *Am J Public Health* 1986;76:585-6.
9. Sorensen G, Pechacek T, Pallonen U. Occupational and worksite norms and attitudes about smoking cessation. *Am J Public Health* 1986;76:544-9.
10. Anonymous. Other people's tobacco smoke [Editorial]. *Lancet* 1986;1:752.
11. Thornberry OT, Wilson RW, Golden PM. Health promotion data for the 1990 objectives. Estimates from the National Health Interview Survey of Health Promotion and Disease Prevention, United States, 1985. In: National Center for Health Statistics. *Advance data from vital and health statistics*. Hyattsville, Maryland: National Center for Health Statistics, 1986; DHHS publication no. (PHS) 86-1250, no. 126.
12. Office on Smoking and Health. *The health consequences of smoking—cancer and chronic lung disease in the workplace: a report of the Surgeon General*. Rockville, Maryland: Public Health Service, 1985. DHHS publication no. (PHS) 85-50207.
13. American Lung Association. Gallup survey [Press release]. New York: American Lung Association, December 5, 1985.

*Epidemiologic Notes and Reports***Tularemia — New Jersey**

On December 3, 1985, a 67-year-old woman died from tularemia in a New Jersey hospital. She had been admitted 7 days previously with a metabolic acidosis secondary to combined dehydration and sepsis. On admission, she had an "unhealing sore" on the first finger of her right hand. Initial treatment included gentamicin and cefazolin, as well as insulin for uncontrolled, late-onset diabetes. After 3 days, the treatment was changed to streptomycin. Despite these measures, disseminated intravascular coagulation, respiratory failure, and hypotension developed, and the woman died.

The case history showed that on November 9, 1985, an 18-year-old neighbor had shot two rabbits behind his home in Gloucester County, New Jersey. After eviscerating the animals, he gave them to the patient and her 64-year-old husband, who skinned and froze the rabbits. During the summer, the young man had noticed several dead rabbits around his house and had attributed their deaths to insecticide that had been sprayed on local fields. One of the two rabbits he shot was noted to be losing its fur.

Two days after dressing out the rabbits, the young man became ill with an ulcerated hand lesion, axillary lymphadenopathy, and a fever. He was examined at the local hospital; no diagnosis was made, but he was treated with antipyretics. On November 23, his two neighbors—the recipients of the rabbits—were admitted to the local hospital. They both had sepsis and hand lesions. On November 26, following instructions from the hospital, the young hunter was started on streptomycin, and he recovered rapidly.

## Tularemia - Continued

The woman's original titer for tularemia, drawn November 23, was less than 20. Her titer rose to 160 after 10 days. First samples from both men were drawn late in the disease. The hunter's first blood specimen was drawn on November 29, when his titer was 1,280. It was reported as 2,560 after 7 days. Blood specimens from the husband were drawn December 3, when his titer was 320, and the level rose to 1,280 after 14 days.

The two rabbits were sent to CDC for analysis. Cultures from the bone marrow of both animals grew *Francisella tularensis*.

Reported by WE Parkin, DVM, State Epidemiologist, New Jersey State Dept of Health; Div of Field Svcs, Epidemiology Program Office, Div of Bacterial Diseases, Center for Infectious Diseases, CDC.

**Editorial Note:** Six cases of tularemia had been reported in New Jersey over the 5-year period prior to this outbreak. One case, in 1985, was also associated with rabbits. No tularemia deaths had been reported in the state in the previous 5 years.

The association between rabbits and human tularemia was first documented in 1913 (7), and rabbit contact was implicated in 90% of the more than 14,000 cases reported through

(Continued on page 753)

TABLE 1. Summary—cases specified notifiable diseases, United States

Disease	48th Week Ending			Cumulative, 48th Week Ending		
	Nov. 29, 1986	Nov. 30, 1985	Median 1981-1985	Nov. 29, 1986	Nov. 30, 1985	Median 1981-1985
Acquired Immunodeficiency Syndrome (AIDS)	78	110	N	12,052	7,280	N
Asplenic meningitis	146	173	222	9,715	9,631	8,956
Encephalitis: Primary (arthropod-borne & unspecified)	10	9	22	1,113	1,220	1,430
Post-infectious	-	1	2	93	112	83
Gonorrhea: Civilian	13,609	14,484	15,535	817,442	819,309	831,394
Military	218	216	240	15,544	19,346	22,245
Hepatitis: Type A	416	518	476	20,729	21,099	21,099
Type B	390	546	512	23,431	24,159	22,014
Non A, Non B	37	59	N	3,182	3,764	N
Unspecified	64	111	125	4,074	5,297	6,686
Legionellosis	11	20	N	733	713	N
Leptospirosis	4	4	4	228	342	221
Malaria	15	12	14	1,027	951	951
Measles: Total*	29	34	15	5,914	2,704	2,516
Indigenous	28	34	N	5,616	2,272	N
Imported	1	-	N	292	432	N
Meningococcal infections: Total	31	40	47	2,214	2,190	2,508
Civilian	31	40	47	2,212	2,183	2,493
Military	-	-	-	2	-	13
Mumps	167	52	67	4,878	2,709	3,044
Pertussis	37	65	36	3,943	3,275	2,150
Rubella (German measles)	3	4	11	477	601	914
Syphilis (Primary & Secondary): Civilian	425	463	521	24,840	24,817	28,595
Military	2	3	3	146	150	351
Toxic Shock syndrome	7	7	N	318	345	N
Tuberculosis	255	334	515	20,103	19,619	21,575
Typhoid fever	-	3	3	150	170	260
Typhus fever, tick-borne (RMSF)	4	4	4	289	354	365
Rabies, animal	2	4	5	735	675	949
	44	68	90	4,392	4,982	5,602

TABLE 2. Notifiable diseases of low frequency, United States

	Cum. 1986		Cum. 1986
Anthrax	-	Leptospirosis (Tex. 1)	37
Botulism: Foodborne	18	Plague	8
Infant (Tex. 1, Utah 1, Oreg. 1)	64	Polioencephalitis, Paralytic	1
Other	1	Psittacosis (Mich. 1)	88
Brucellosis	77	Rabies, human	-
Cholera	16	Tetanus (Tex. 1)	59
Congenital rubella syndrome	10	Trichinosis	31
Congenital syphilis, ages < 1 year	107	Typhus fever, flea-borne (endemic, murine)	46
Diphtheria	-		

\*One of the 29 reported cases for this week was imported from a foreign country or can be directly traceable to a known internationally imported case within two generations.

TABLE III. Cases of specified notifiable diseases, United States, weeks ending  
November 29, 1986 and November 30, 1985 (48th Week)

Reporting Area	AIDS Cum 1986	Aseptic Mening- itis 1986	Encephalitis		Gonorrhea (Civilian)		Hepatitis (Viral, by type)				Legionel- losis 1986	Leprosy Cum 1986
			Primary Cum 1986	Post-in- fectious Cum 1986	Cum 1986	Cum 1985	A 1986	B 1986	NA,NB 1986	Unspeci- fied 1986		
UNITED STATES	12,052	146	1,113	93	817,442	819,309	416	390	37	64	11	228
NEW ENGLAND	479	10	29	3	22,013	20,879	12	41	-	11	1	8
Maine	20	-	2	-	789	1,067	1	1	-	-	-	-
N.H.	13	-	-	-	529	523	-	-	-	-	-	-
VT	5	2	4	2	246	307	-	-	-	1	1	-
Mass	254	1	5	-	7,983	8,636	2	30	-	10	-	8
RI	31	3	-	-	1,710	1,724	3	3	-	-	-	-
Conn	156	4	16	1	10,756	8,622	6	7	-	-	-	-
MID ATLANTIC	4,417	8	97	10	142,792	119,042	7	23	-	21	-	17
Upstate N.Y.	487	3	35	6	17,218	16,772	3	8	-	-	-	1
N.Y. City	2,998	3	19	1	82,811	58,145	-	-	-	19	-	15
N.J.	664	2	10	-	18,120	17,856	4	15	-	2	-	-
Pa.	268	-	33	3	24,643	26,289	-	-	-	-	-	1
E.N. CENTRAL	725	19	338	11	106,249	107,001	5	30	-	1	3	4
Ohio	154	9	130	3	27,344	29,345	2	22	-	-	3	-
Ind	59	7	79	3	11,380	11,479	1	2	-	1	-	-
Ill	347	-	50	4	25,340	25,109	-	-	-	-	-	4
Mich	127	3	53	1	34,685	30,751	2	6	-	-	-	1
Wis	38	-	26	-	7,248	10,317	-	-	-	-	-	-
W.N. CENTRAL	225	14	81	9	35,215	38,202	12	13	2	-	-	4
Minn	88	-	34	-	5,064	5,649	6	5	-	-	-	2
Iowa	18	2	26	-	3,602	4,114	-	-	-	-	-	-
Mo	72	9	2	-	17,429	18,423	1	8	1	-	-	-
N.Dak	2	-	4	-	290	258	-	-	-	-	-	-
S.Dak	2	-	11	-	720	736	2	-	1	-	-	-
Nebr	10	-	1	-	2,802	3,257	1	-	-	-	-	-
Kans	33	3	3	8	5,488	5,765	2	-	-	-	-	2
S ATLANTIC	1,735	17	144	37	211,254	214,043	32	113	13	5	2	3
Del	22	-	6	-	3,483	4,152	-	-	-	-	-	-
Md	183	3	31	1	25,081	27,216	-	2	1	-	-	-
D.C.	222	1	-	1	15,870	14,663	2	2	-	-	-	-
Va	137	2	40	1	17,520	17,866	4	7	2	1	2	1
W.Va	7	1	45	-	2,063	2,419	-	4	1	-	-	-
N.C.	73	4	18	2	32,829	33,800	1	8	3	1	-	-
S.C.	47	-	-	-	17,943	20,199	6	22	-	-	-	-
Ge	266	2	-	1	35,232	41,607	6	21	-	-	-	-
Fla	778	4	4	31	61,243	52,121	12	47	6	3	-	2
E.S. CENTRAL	147	20	61	4	65,474	70,921	2	20	-	-	-	1
Ky	28	11	30	1	7,214	8,111	-	1	-	-	-	-
Tenn	70	2	8	1	24,787	27,279	1	10	-	-	-	-
Ala	25	7	22	2	19,225	21,251	1	8	-	-	-	1
Miss	24	-	1	-	14,248	14,280	-	1	-	-	-	-
W.S. CENTRAL	1,068	28	180	6	95,304	103,378	39	23	2	9	1	23
Ark	29	-	-	2	9,018	9,666	1	2	-	-	-	1
La	143	3	16	-	16,399	19,430	3	4	-	-	1	1
Okla	41	2	21	-	10,917	11,547	5	3	1	-	-	-
Tex	855	23	143	4	58,970	62,735	30	14	1	9	-	21
MOUNTAIN	323	5	38	1	24,150	25,954	77	37	2	5	1	13
Mont	4	-	1	1	833	746	1	1	-	-	1	-
Idaho	3	-	-	-	800	897	8	2	-	-	-	-
Wyo	4	-	2	-	800	594	-	-	-	-	-	-
Calo	156	1	5	-	6,208	7,553	4	5	-	2	-	3
N.Mex	23	-	3	-	2,556	2,878	31	7	1	-	-	-
Ariz	80	4	18	-	7,783	7,911	32	18	-	3	-	7
Utah	18	-	7	-	1,029	1,259	-	3	-	-	-	1
Nev	35	-	2	-	4,641	4,116	1	1	1	-	-	2
PACIFIC	2,933	25	145	12	114,991	119,889	230	90	18	12	3	155
Wash	157	2	13	-	8,367	9,246	46	14	4	3	-	16
Oreg	58	-	-	-	5,061	5,950	25	8	4	-	-	-
Calif	2,656	21	124	12	98,190	100,250	151	68	9	8	3	105
Alaska	12	-	7	-	2,442	2,870	8	-	1	1	-	-
Hawaii	50	2	1	-	1,193	1,573	-	-	-	-	-	33
Guam	-	-	-	-	201	181	-	-	-	-	-	1
P.R.	139	-	5	1	2,237	2,912	-	1	-	-	-	7
V.I.	5	-	-	-	254	378	-	1	-	-	-	-
Pac. Trust Terr	-	-	-	-	429	766	-	-	-	-	-	56
Amer Samoa	-	-	-	-	53	-	-	-	-	-	-	3

N Not notifiable

U Unavailable

TABLE III. (Cont'd.) Cases of specified notifiable diseases, United States, weeks ending  
November 29, 1986 and November 30, 1985 (48th Week)

Reporting Area	Malina	Measles (Rubella)				Meningo- coccal infections	Mumps		Pertussis			Rubella			
		Indigenous		Imported *			Total	Cum 1986	Cum 1986	1986	Cum 1986	Cum 1985	1986	Cum 1986	Cum 1985
		1986	Cum 1986	1986	Cum 1986										
UNITED STATES	1,027	28	5,616	1	292	2,704	2,214	167	4,878	37	3,943	3,275	3	477	601
NEW ENGLAND	62	-	88	-	16	128	159	-	65	2	189	204	-	9	13
Maine	2	-	12	-	1	1	27	-	-	-	2	9	-	-	-
NH	3	-	43	-	-	-	6	-	14	1	82	112	-	1	3
Vt	2	-	-	-	-	-	19	-	4	-	3	3	-	-	-
Mass	32	-	24	-	13	118	45	-	14	1	42	47	-	4	6
RI	7	-	2	-	-	-	21	-	11	-	6	22	-	2	-
Conn	16	-	7	-	2	7	41	-	22	-	24	11	-	1	4
MID ATLANTIC	141	-	1,729	-	34	232	344	5	203	5	202	242	-	37	228
Upstate N.Y.	48	-	77	-	24	85	122	1	66	1	125	115	-	27	18
N.Y. City	31	-	723	-	4	79	70	-	29	-	10	29	-	5	185
N.J.	37	-	906	-	4	28	30	1	51	2	20	11	-	5	11
Pa.	25	-	24	-	2	40	122	3	57	2	47	87	-	-	14
E.N. CENTRAL	61	-	1,079	-	28	582	321	89	3,173	8	375	799	-	48	38
Ohio	19	-	-	-	10	60	132	4	132	8	167	117	-	1	3
Ind	2	-	27	-	11	57	34	-	40	-	35	201	-	-	-
Ill	16	-	899	-	4	346	74	84	2,412	-	36	74	-	38	20
Mich	20	-	75	-	-	60	69	1	339	-	35	48	-	8	16
Wis	4	-	278	-	3	59	11	-	250	-	99	359	-	2	1
W.N. CENTRAL	30	-	323	-	17	12	104	12	161	-	1,406	234	-	14	19
Miss	8	-	45	-	4	6	23	5	18	-	48	116	-	1	2
Iowa	1	-	133	-	1	-	11	7	57	-	19	31	-	1	1
Mo	12	-	26	-	6	3	34	-	24	-	22	32	-	1	7
N. Dak	-	-	25	-	1	2	1	-	4	-	5	10	-	-	-
S. Dak	3	-	-	-	-	-	5	-	1	-	14	5	-	-	-
Nebr	4	-	-	-	-	-	11	-	-	-	10	11	-	-	-
Kans	3	-	94	-	5	1	19	-	57	-	1,288	29	-	10	7
S. ATLANTIC	122	27	775	-	56	340	403	7	242	5	749	535	2	11	52
Del	1	-	1	-	-	-	5	-	1	-	227	2	-	-	2
Md	14	-	26	-	9	115	46	1	26	-	164	311	-	-	6
D.C.	6	-	-	-	1	31	5	-	1	-	-	-	-	-	-
Va	32	-	36	-	24	28	71	1	44	-	41	20	-	-	2
W. Va	4	-	2	-	3	4	1	-	1	-	26	4	-	-	9
N.C.	6	-	3	-	1	9	63	1	28	3	79	34	-	-	1
S.C.	6	-	274	-	-	3	44	1	15	-	18	2	-	-	3
Ga	13	-	79	-	14	8	58	-	28	-	132	92	-	-	-
Fla	40	27	354	-	7	113	107	3	51	1	62	70	2	11	29
E.S. CENTRAL	21	-	63	-	9	7	114	39	164	-	47	67	-	4	3
Ky	6	-	-	-	6	5	26	-	6	-	16	26	-	-	-
Tenn	1	-	57	-	1	1	37	39	153	-	10	28	-	-	-
Ala	10	-	1	-	1	-	37	-	4	-	25	28	-	-	-
Miss	4	-	5	-	1	1	14	-	1	-	1	7	-	-	-
W.S. CENTRAL	103	-	680	-	38	440	206	3	268	-	250	533	-	71	42
Ark	1	-	276	-	2	-	27	-	61	-	20	14	-	1	1
La	18	-	4	-	-	42	26	-	3	-	15	17	-	-	-
Okla	12	-	37	-	2	1	30	N	N	-	126	166	-	-	2
Tex	72	-	363	-	34	397	123	3	204	-	89	336	-	70	39
MOUNTAIN	38	-	302	-	29	541	104	3	248	4	273	220	-	24	6
Mont	1	-	-	-	8	137	10	1	6	-	20	10	-	2	-
Idaho	1	-	1	-	-	137	4	-	8	4	46	17	-	-	2
Wyo	-	-	-	-	-	5	2	-	-	-	4	1	-	-	-
Colo	12	-	2	-	8	15	19	1	17	-	66	87	-	1	-
N. Mex	5	-	33	-	7	6	11	N	N	-	26	12	-	-	2
Ariz	13	-	252	-	6	241	22	-	193	-	85	40	-	2	1
Utah	3	-	12	-	-	-	10	-	15	-	42	53	-	15	-
Nev	3	-	2	-	-	-	26	1	9	-	4	-	-	3	1
PACIFIC	449	1	577	1	85	424	459	9	354	13	482	441	1	258	200
Wash	29	1	140	-	28	142	63	-	18	-	149	82	-	17	14
Oreg	19	-	7	-	4	5	36	N	14	2	50	-	-	4	2
Calif	400	-	403	1	31	253	334	7	305	6	297	262	1	231	136
Alaska	-	-	-	-	-	-	14	2	8	1	5	30	-	-	1
Hawaii	1	-	27	-	2	24	13	-	23	4	20	17	-	6	48
Guam	1	-	4	-	1	11	1	-	4	-	-	-	-	4	3
P.R.	4	-	36	-	-	67	3	-	33	-	19	15	-	62	27
V.I.	-	-	-	-	-	10	-	-	17	-	-	-	-	-	-
Pac. Trust Terr	-	-	-	-	-	-	1	-	11	-	-	-	-	2	-
Amer Samoa	-	-	2	-	-	-	-	-	5	-	-	-	-	1	-

\*For measles only, imported cases includes both out-of-state and international importations.

N Not notifiable

U Unavailable

I International

O Out-of-state

TABLE III. (Cont'd.) Cases of specified notifiable diseases, United States, weeks ending  
November 29, 1986 and November 30, 1985 (48th Week)

Reporting Area	Syphilis (Civilian) (Primary & Secondary)		Toxic- shock Syndrome	Tuberculosis		Tula- remia	Typhoid Fever	Typhus Fever (Tick-borne) (RMSF)	Rabies, Animal
	Cum 1986	Cum 1985		Cum 1986	Cum 1985				
UNITED STATES	24,840	24,817	7	20,103	19,819	150	289	735	4,992
NEW ENGLAND	457	547	-	625	664	1	16	13	8
Maine	19	15	-	34	45	-	-	-	-
NH	10	38	-	23	21	-	-	2	1
VT	9	7	-	16	8	-	-	-	2
Mass	246	270	-	347	391	1	13	4	-
RI	19	17	-	42	50	-	-	3	3
Conn	154	200	-	163	149	-	3	4	2
MID ATLANTIC	3,492	3,340	3	3,974	3,502	1	24	40	634
Upstate N.Y.	170	244	-	576	600	-	4	20	81
N.Y. City	1,958	2,030	-	2,078	1,715	-	11	5	-
N.J.	609	643	3	673	479	1	8	2	17
Pa.	755	431	-	647	708	-	1	13	536
E.N. CENTRAL	792	917	-	2,379	2,394	1	23	46	134
Ohio	117	135	-	419	410	-	9	40	16
Ind	103	75	-	258	309	-	2	-	17
Ill	370	414	-	1,026	1,042	-	3	2	39
Mich	162	233	-	572	498	1	6	4	24
Wis	40	60	-	104	135	-	3	-	38
W.N. CENTRAL	198	218	-	585	580	41	9	48	766
Minn	31	42	-	136	118	-	2	1	120
Iowa	8	16	-	46	53	1	-	1	177
Mo	104	121	-	289	270	30	6	24	67
N. Dak	5	2	-	10	10	-	-	1	146
S. Dak	9	6	-	28	31	3	-	6	170
Nebr	11	7	-	14	16	1	-	5	32
Kans	30	22	-	62	62	6	1	10	54
S. ATLANTIC	7,455	7,113	-	4,048	4,031	12	45	330	1,263
Del	53	36	-	42	42	-	1	1	1
Md	423	461	-	289	360	2	15	29	559
D.C.	274	306	-	152	141	1	4	-	31
Va	316	282	-	344	406	3	10	51	189
W. Va	20	25	-	115	102	-	3	10	53
N.C.	488	634	-	596	538	3	4	128	10
S.C.	646	743	-	516	489	-	-	70	64
Ga	1,391	1,287	-	668	677	3	-	39	193
Fla	3,844	3,339	-	1,326	1,276	-	8	2	163
E.S. CENTRAL	1,867	1,933	-	1,772	1,692	13	4	110	354
Ky	65	65	-	404	414	5	-	22	100
Tenn	575	592	-	508	502	6	1	45	138
Ala	485	611	-	557	499	1	1	25	113
Miss	542	665	-	303	277	1	2	18	3
W.S. CENTRAL	4,867	5,743	1	2,541	2,508	67	29	137	685
Ark	243	308	-	343	292	49	-	16	155
La	847	1,008	-	393	369	1	1	1	22
Okla	137	178	-	235	236	12	2	103	57
Tex	3,640	4,249	1	1,570	1,611	5	26	17	451
MOUNTAIN	561	714	-	494	527	11	16	10	630
Mont	7	6	-	31	46	1	1	4	204
Idaho	14	7	-	23	25	-	-	2	9
Wyo	4	14	-	-	7	1	-	1	266
Colo	126	201	-	51	83	3	1	3	29
N. Mex	68	120	-	92	83	1	1	-	6
Ariz	233	297	-	231	230	-	9	-	98
Utah	18	9	-	31	17	4	3	-	7
Nev	91	60	-	35	36	1	1	-	11
PACIFIC	5,351	4,284	3	3,685	3,741	3	123	1	518
Wash	152	99	1	202	204	1	3	-	5
Oreg	107	100	-	118	124	-	-	-	1
Calif	5,047	4,014	2	3,154	3,141	1	114	1	504
Alaska	10	4	-	46	95	1	1	-	8
Hawaii	35	67	-	165	177	-	5	-	-
Guam	1	2	-	34	38	-	1	-	-
P.R.	808	813	-	305	330	-	5	-	43
VI	1	3	-	1	-	-	-	-	-
Pac. Trust Terr	246	128	-	81	75	-	49	-	-
Amer Samoa	-	-	-	5	-	-	-	-	-

U Unavailable



TABLE IV. Deaths in 121 U.S. cities.\* week ending  
November 29, 1986 (48th Week)

Reporting Area	All Causes, By Age (Years)						P&T <sup>††</sup> Total	Reporting Area	All Causes, By Age (Years)						P&T <sup>††</sup> Total
	All Ages	≥85	45-64	25-44	1-24	<1			All Ages	≥85	45-64	25-44	1-24	<1	
NEW ENGLAND	552	397	98	31	8	18	42	S ATLANTIC	971	582	245	89	24	29	37
Boston, Mass.	118	68	30	7	4	9	14	Atlanta, Ga.	95	65	22	5	2	1	2
Bridgeport, Conn.	40	29	8	3	-	1	1	Baltimore, Md.	184	81	51	19	5	8	10
Cambridge, Mass.	29	22	4	3	-	-	3	Charlotte, N.C.	61	40	12	6	3	-	1
Fall River, Mass.	18	14	4	-	-	-	-	Jacksonville, Fla.	58	41	15	2	-	-	1
Hartford, Conn.	64	46	11	4	2	1	2	Miami, Fla.	86	55	19	8	2	2	2
Lowell, Mass.	18	13	2	2	1	-	-	Norfolk, Va.	52	37	13	2	-	-	1
Lynn, Mass.	17	17	-	-	-	-	-	Richmond, Va.	53	27	20	5	-	1	2
New Bedford, Mass.	19	16	2	1	-	-	1	Savannah, Ga.	24	12	8	2	-	2	2
New Haven, Conn.	50	30	10	7	-	3	5	St. Petersburg, Fla.	86	67	12	6	1	-	4
Providence, R.I.	32	27	3	1	-	1	2	Tampa, Fla.	42	20	9	7	2	2	3
Somerville, Mass.	9	5	4	-	-	-	1	Washington, D.C.	218	110	59	27	9	13	8
Springfield, Mass.	42	38	3	1	-	-	4	Wilmington, Del.	32	27	5	-	-	-	1
Waterbury, Conn.	31	24	4	2	-	-	1								
Worcester, Mass.	65	48	13	-	1	3	3								
MID ATLANTIC	2,617	1,714	531	263	56	53	121	E.S. CENTRAL	635	392	155	48	21	19	25
Albany, N.Y.	50	32	7	7	3	1	2	Birmingham, Ala.	101	80	28	11	3	1	1
Albion, N.Y.	23	18	5	-	-	-	1	Chattanooga, Tenn.	39	26	10	2	-	1	2
Albion, Pa.	77	49	18	9	2	1	8	Knoxville, Tenn.	77	47	17	6	3	4	2
Buffalo, N.Y.	25	15	4	4	1	1	2	Louisville, Ky.	122	78	31	8	3	2	8
Camden, N.J.	28	20	4	4	-	-	2	Memphis, Tenn.	120	68	30	8	9	7	4
Elizabeth, N.J.	37	27	6	1	1	2	4	Mobile, Ala.	38	26	6	2	2	2	2
Encl. Pa.	37	26	9	2	-	-	4	Montgomery, Ala.	29	22	5	2	-	-	1
Jersey City, N.J.	1,483	923	396	193	34	27	43	Nashville, Tenn.	109	67	30	9	1	2	5
N.Y. City, N.Y.	40	25	9	5	-	-	1								
Newark, N.J.	38	12	10	7	2	7	3	W.S. CENTRAL	968	563	242	98	33	32	32
Pateron, N.J.	371	244	89	26	6	6	24	Austin, Tex.	47	33	10	3	1	-	1
Philadelphia, Pa.	72	47	17	6	1	1	3	Baton Rouge, La.	17	8	6	1	2	-	-
Pittsburgh, Pa.	31	24	5	2	-	-	2	Corpus Christi, Tex.	30	18	8	3	-	1	1
Reading, Pa.	114	86	16	4	4	2	10	Dallas, Tex.	154	81	47	21	4	1	2
Rochester, N.Y.	23	16	7	-	-	-	1	El Paso, Tex.	41	25	11	3	-	2	4
Schenectady, N.Y.	22	20	2	-	-	-	1	Fort Worth, Tex.	60	34	15	3	4	4	3
Scranton, Pa.	90	62	19	6	2	1	5	Houston, Tex.	304	163	75	39	14	13	7
Syracuse, N.Y.	36	23	7	5	-	-	1	Little Rock, Ark.	51	33	10	4	2	2	4
Trenton, N.J.	13	9	-	2	-	-	2	New Orleans, La.	80	50	16	9	1	4	-
Utica, N.Y.	37	34	3	-	-	-	8	San Antonio, Tex.	131	83	30	11	3	4	9
Yonkers, N.Y.	1,938	1,271	408	145	47	64	82	Shreveport, La.	5	5	-	-	-	-	-
								Tulsa, Okla.	48	30	14	1	2	1	1
E.N. CENTRAL	1,938	1,271	408	145	47	64	82	MOUNTAIN	517	331	114	37	17	18	18
Akron, Ohio	28	24	3	2	-	-	-	Albuquerque, N.Mex.	47	31	3	9	2	2	1
Canton, Ohio	28	21	5	1	-	-	1	Colorado Springs, Colo.	31	21	2	2	3	3	1
Chicago, Ill.	564	362	125	45	10	22	16	Denver, Colo.	99	60	31	5	2	1	4
Cincinnati, Ohio	138	93	29	8	3	5	9	Las Vegas, Nev.	73	43	23	4	3	-	2
Cleveland, Ohio	143	82	37	13	1	10	1	Ogden, Utah	17	10	5	-	1	1	-
Columbus, Ohio	175	104	44	16	7	4	12	Phoenix, Ariz.	126	78	28	10	4	6	5
Dayton, Ohio	76	48	18	6	1	2	2	Pueblo, Colo.	21	16	3	2	-	-	3
Detroit, Mich.	150	89	26	19	9	7	4	Salt Lake City, Utah	39	25	8	1	2	3	-
Evansville, Ind.	19	13	3	-	1	2	1	Tucson, Ariz.	64	47	11	4	-	2	2
Fort Wayne, Ind.	53	34	9	7	2	1	4								
Gary, Ind.	13	6	4	2	1	-	1	PACIFIC	1,491	1,007	266	124	41	46	92
Grand Rapids, Mich.	27	21	3	-	2	1	1	Berkeley, Calif.	13	10	2	-	-	1	1
Indianapolis, Ind.	33	90	30	8	3	1	4	Fresno, Calif.	63	46	7	4	2	2	4
Madison, Wis.	48	23	7	1	1	4	1	Glendale, Calif.	22	17	4	1	-	-	1
Milwaukee, Wis.	97	73	18	3	1	2	5	Honolulu, Hawaii	49	27	14	4	2	2	8
Peoria, Ill.	31	25	3	-	1	2	4	Long Beach, Calif.	108	67	25	12	3	1	12
Rockford, Ill.	49	38	9	2	-	-	3	Los Angeles, Calif.	302	208	51	22	12	4	8
South Bend, Ind.	41	27	8	4	2	-	4	Oakland, Calif.	59	42	10	4	2	1	2
Toledo, Ohio	73	58	11	1	1	2	2	Pasadena, Calif.	25	19	5	1	-	-	-
Youngstown, Ohio	62	40	16	4	1	1	1	Portland, Oreg.	144	100	25	13	4	2	9
								Sacramento, Calif.	97	66	17	7	3	4	11
W.N. CENTRAL	758	520	143	52	22	21	31	San Diego, Calif.	114	71	24	12	2	4	6
Des Moines, Iowa	56	43	11	2	-	-	3	San Francisco, Calif.	141	85	29	19	1	6	4
Duluth, Minn.	12	14	2	1	-	-	-	San Jose, Calif.	139	82	29	15	3	10	13
Kansas City, Kans.	23	16	2	1	3	1	-	Seattle, Wash.	110	110	19	8	7	6	7
Kansas City, Mo.	131	88	25	10	3	5	7	Spokane, Wash.	44	40	2	-	1	1	7
Lincoln, Neb.	27	22	4	-	-	1	1	Tacoma, Wash.	21	15	3	2	-	1	-
Minneapolis, Minn.	184	133	30	11	4	6	5								
Omaha, Neb.	58	35	12	6	2	3	2								
St. Louis, Mo.	148	90	34	17	4	3	6								
St. Paul, Minn.	60	46	9	-	4	1	2								
Wichita, Kans.	54	33	14	4	2	1	5								
TOTAL	10,444 <sup>††</sup>	6,777	2,202	887	269	300	480								

\* Mortality data in this table are voluntarily reported from 121 cities in the United States, most of which have populations of 100,000 or more. A death is reported by the place of its occurrence and by the week that the death certificate was filed. Fetal deaths are not included.

<sup>††</sup> Pneumonia and influenza.

<sup>†††</sup> Because of changes in reporting methods in these 3 Pennsylvania cities, these numbers are partial counts for the current week. Complete counts will be available in 4 to 6 weeks.

<sup>††††</sup> Total includes unknown ages.

<sup>†††††</sup> Data not available. Figures are estimates based on average of past 4 weeks.



### Tularemia — Continued

1944 (2). Although bloodsucking anthropod vectors have accounted for an increasing percentage of cases in more recent years, rabbits continue to be an important source of infection (3).

In the United States, wild rabbits of the genus *Sylvilagus* (cottontails, marsh rabbits, and swamp rabbits) present the greatest hazard (2). Jack rabbits and snowshoe hares are susceptible to tularemia but have rarely been implicated as direct sources of human infection. The domestic rabbit (*Oryctolagus cuniculus*) has not been documented as a source of human tularemia (2).

Exposure of the skin or conjunctiva to blood and other infectious tissue while skinning and dressing rabbits account for most cases. Ingestion of inadequately cooked meat has also been implicated. In some instances pulmonary tularemia has resulted from breathing aerosols generated while handling unopened rabbits or merely by poking at a dead rabbit with a stick (4,5). Indirect transmission from rabbits to humans may result from bites by pet animals or deerflies (6,7).

In 1939, the peak incidence year in the United States, 2,291 cases (17.5/1,000,000 population) were reported (2). Only 291 cases (1.2/1,000,000 population) were reported in 1984.

### References

1. Wherry WB, Lamb BH. Infection of man with *Bacterium tularensis*. J Infect Dis 1914;15:331-40.
2. Jellison WL. Tularemia in North America 1930-1974. Missoula, Montana: University of Montana, 1974.
3. Boyce JM. Recent trends in the epidemiology of tularemia in the United States. J Infect Dis 1975;131:197-9.
4. Martone WJ, Marshall LW, Kaufmann AF, Hobbs JH, Levy ME. Tularemia pneumonia in Washington, DC. A report of three cases with possible common-source exposures. JAMA 1979;242:2315-7.
5. Halsted CC, Kulasinghe HP. Tularemia pneumonia in urban children. Pediatrics 1978;61:860-2.
6. Evans ME, McGee ZA, Hunter PT, Schaffner W. Tularemia and the tomcat. JAMA 1981; 246:1343.
7. Klock LE, Olsen PF, Fukushima T. Tularemia epidemic associated with the deerfly. JAMA 1973; 226:149-52.

### Multiply Resistant Shigellosis in a Day-Care Center — Texas

Between October 10 and November 6, 1985, 15 children at a day-care center in Diboll, Texas, developed a diarrheal illness. *Shigella sonnei* was isolated from 10 ill children and from two of 19 asymptomatic children who were cultured on November 7. All isolates were colicin type 9, resistant to ampicillin, carbenicillin, streptomycin, cephalothin, and trimethoprim/sulfamethoxazole (TMP/SMX), and sensitive to tetracycline, nalidixic acid, chloramphenicol, and gentamicin. The attack rate was highest among the 12- to 22-month-old group. Family members of this group had the highest secondary attack rate (Table 1). No cases occurred among the 22 staff members. None of the children were hospitalized, but four of the five ill family members were.

The 89 children attending the center were cared for, by age group, in separate rooms. All groups except infants and toddlers had separate toilet and playground facilities. Infants and toddlers shared these facilities.

*Shigellosis — Continued*

Symptomatic children were excluded from the center until their diarrhea had resolved. Then they were permitted to return, without treatment or cultures, to their classrooms. Handwashing and hygiene were emphasized; contact between age groups was limited; and the routine policy excluding food preparers from child care, particularly diaper-changing, was reinforced. No further cases were reported at the center after November 7, when this strategy was implemented.

During the following month, statewide surveillance for TMP/SMX-resistant *S. sonnei* infections detected an outbreak among kindergarteners in a town 100 miles away. Although this outbreak strain had the same colicin type and antimicrobial resistance profile as the Diboll strain, its plasmid content differed, and no direct connection between the two outbreaks was discovered.

Reported by M Crowder, MD, W Joyce, J Connors, Public Health Region 7, A Quillion, M Czpiel, Angelina County and Cities Health Dist, J Taylor, MPH, DL Martin, MN, CE Alexander, MD, Bureau of Epidemiology, Texas Dept of Health; Enteric Diseases Br, Div of Bacterial Diseases, Center for Infectious Diseases, CDC.

**Editorial Note:** Shigellosis in day-care centers can be difficult to control. Basic hygiene, exclusion of symptomatic persons, and routine antimicrobial therapy for all infected persons have been advocated as control measures (1). In the Texas outbreak reported here, antimicrobial therapy was not part of the control strategy because the strain was resistant to all drugs commonly used to treat shigellosis in children. Nonetheless, the straightforward control strategy in this well-designed day-care center was associated with the end of the outbreak, even though untreated convalescent children returned to the center and untreated asymptomatic carrier children remained there.

The elements contributing to this apparent success included vigorous emphasis on handwashing among staff and children; routine exclusion of ill children; separate areas and staff for diapering and food-preparation; and separate rooms, toilets, and play-facilities for different age groups. There is some evidence that each element is important. Handwashing has been shown to reduce the incidence of diarrheal illness in day-care centers (2). In day-care centers in Houston, Texas, the incidence of diarrheal illness was significantly associated with the proportion of staff who changed diapers and also served or prepared food (3). The usefulness of separating children by age was suggested by uniform shigellosis attack rates observed across ages 0 to 5 years at a day-care center where the children were grouped together (4). Additional study of the efficacy and utility of these specific control measures is needed (5,6).

Providing day-care in isolation for convalescent children may limit the spread of shigellosis in the community. In one outbreak, in which children with shigellosis were rigidly excluded from a day-care center until negative cultures were obtained, the outbreak strain spread to a day-care center in an adjacent county (7). In another outbreak, at a center where isolation of

TABLE 1. Attack rates of shigellosis, by room assignment, among children < 5 years old attending a day-care center and their family members — Diboll, Texas, 1985.

Room assignment	Cases/Children	Attack rate (%)	Cases/Family members	Attack rate (%)
Infant	1/8	12	0/20	0
Toddler	6/10	60	4/31	13
Two-year-old	3/15	20	1/40	2
Three-year-old	4/10	40	0/29	0
Four-year-old	1/15	7	0/47	0
Total	15/58	26	5/167	3

*Shigellosis — Continued*

convalescent children was possible, treated, convalescent children without negative cultures were allowed to return to the day-care center, and there was no further spread of illness in either the center or the community (8). Further evaluation of convalescent day-care, with and without isolation, is needed before specific recommendations can be made.

To help day-care center directors, employees, and parents work with health departments to control disease in day-care centers, CDC has produced a training kit: "What To Do To Stop Disease in Child Day-Care Centers". This kit has been distributed to state health departments and licensing boards for distribution to licensed day-care centers. It also can be purchased for \$4.00 from the Government Printing Office, Superintendent of Documents, Washington, D.C., 20402. The GPO Stock Number is 017-023-00172-8.

*References*


1. Child Day Care Infectious Disease Study Group, CDC. Public health considerations of infectious diseases in child day care centers. *J Pediatr* 1984;105:683-701.
2. Black RE, Dykes AC, Anderson KE, et al. Handwashing to prevent diarrhea in day-care centers. *Am J Epidemiol* 1981;113:445-51.
3. Lemp GF, Woodward WE, Pickering LK, Sullivan PS, DuPont HL. The relationship of staff to the incidence of diarrhea in day-care centers. *Am J Epidemiol* 1984;120:750-8.
4. Pickering LK, Evans DG, DuPont HL, Vollet JJ 3rd, Evans DJ Jr. Diarrhea caused by *Shigella*, rotavirus and *Giardia* in day-care centers: prospective study. *J Pediatr* 1981;99:51-6.
5. Pickering LK, Bartlett AV, Woodward WE. Acute infectious diarrhea among children in day care: epidemiology and control. *Rev Inf Dis* 1986;8:539-47.
6. Petersen NJ, Bressler GK. Design and modification of the day care environment. *Rev Inf Dis* 1986;8:618-21.
7. Tacket CO, Cohen ML. Shigellosis in day care centers: use of plasmid analysis to assess control measures. *Pediatr Infect Dis* 1983;2:127-30.
8. Tauxe RV, Johnson KE, Boase JC, Helgeson SD, Blake PA. Control of day care shigellosis: a trial of convalescent day care in isolation. *Am J Publ Health* 1986;76:627-30.

**Erratum: Vol. 35, No. 46**

- p. 714 The following reference replaces reference number 5 for the article entitled "Premature Mortality from Diabetes Mellitus—Use of Sentinel Health Event Surveillance to Assess Causes":

5. Tunbridge WM. Factors contributing to deaths of diabetics under fifty years of age. On behalf of the Medical Services Study Group and British Diabetic Association. *Lancet* 1981;2:569-72.

FIGURE I. Reported measles cases — United States, weeks 44-47, 1986



The *Morbidity and Mortality Weekly Report* is prepared by the Centers for Disease Control, Atlanta, Georgia, and available on a paid subscription basis from the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402. (202) 783-3238.

The data in this report are provisional, based on weekly reports to CDC by state health departments. The reporting week concludes at close of business on Friday; compiled data on a national basis are officially released to the public on the succeeding Friday.

The editor welcomes accounts of interesting cases, outbreaks, environmental hazards, or other public health problems of current interest to health officials. Such reports and any other matters pertaining to editorial or other textual considerations should be addressed to: ATTN: Editor, *Morbidity and Mortality Weekly Report*, Centers for Disease Control, Atlanta, Georgia 30333.

Director, Centers for Disease Control  
James O. Mason, M.D., Dr.P.H.  
Director, Epidemiology Program Office  
Carl W. Tyler, Jr., M.D.

Editor  
Michael B. Gregg, M.D.

U.S. Government Printing Office: 1987-730-145/40037 Region IV

DEPARTMENT OF  
HEALTH & HUMAN SERVICES  
Public Health Service  
Centers for Disease Control  
Atlanta GA 30333

Official Business  
Penalty for Private Use \$300



Postage and Fees Paid  
U.S. Dept. of H.H.S.  
HH5 396

A 48106SER 06 8639 9  
SERIALS ACQUISITION DEPT  
UNIVERSITY MICROFILMS  
300 NORTH ZEEB ROAD  
ANN ARBOR, MI 48106

X

